


4. The area enclosed between the curves  $y = x|x|$  and  $y = x - |x|$  is :

- (1)  $\frac{8}{3}$  (2)  $\frac{2}{3}$   
 (3) 1 (4)  $\frac{4}{3}$

Ans. (4)

Sol.



$$A = \int_{-2}^0 -x^2 - 2x = \frac{4}{3}$$

5. 60 words can be made using all the letters of the word BHBJO, with or without meaning. If these words are written as in a dictionary, then the 50<sup>th</sup> word is :

- (1) OBBHJ (2) HBBJO  
 (3) OBBJH (4) JBBOH

Ans. (3)

Sol. B B H J O

[B] \_\_\_\_\_ 4! = 24

[H] \_\_\_\_\_  $\frac{4!}{2!} = 12$

[J] \_\_\_\_\_  $\frac{4!}{2!} = 12$

O B B H J

O B B J H  $\rightarrow$  50<sup>th</sup> rank

6. Let  $\vec{a} = 2\hat{i} + 5\hat{j} - \hat{k}$ ,  $\vec{b} = 2\hat{i} - 2\hat{j} + 2\hat{k}$  and  $\vec{c}$  be three vectors such that  $(\vec{c} + \hat{i}) \times (\vec{a} + \vec{b} + \hat{i}) = \vec{a} \times (\vec{c} + \hat{i})$ .  $\vec{a} \cdot \vec{c} = -29$ , then  $\vec{c} \cdot (-2\hat{i} + \hat{j} + \hat{k})$  is equal to :

- (1) 10 (2) 5  
 (3) 15 (4) 12

Ans. (2)

Sol. Let's assume  $\vec{v} = \vec{a} + \vec{b} + \hat{i}$

$$= 5\hat{i} + 3\hat{j} + \hat{k}$$

and  $\vec{c} + \hat{i} = \vec{p}$

So,

$$\vec{p} \times \vec{v} = \vec{a} \times \vec{p}$$

$$\vec{p} \times \vec{v} + \vec{p} \times \vec{a} = \vec{0}$$

$$\vec{p} \times (\vec{v} + \vec{a}) = \vec{0}$$

$$\Rightarrow \vec{p} = \lambda(\vec{v} + \vec{a})$$

$$\vec{c} + \hat{i} = \lambda(7\hat{i} + 8\hat{j})$$

$$\vec{a} \cdot \vec{c} + \vec{a} \cdot \hat{i} = \lambda \vec{a} \cdot (7\hat{i} + 8\hat{j})$$

$$-29 + 2 = \lambda(14 + 40)$$

$$\lambda = -\frac{1}{2}$$

$$\vec{c} \cdot (-2\hat{i} + \hat{j} + \hat{k}) + \hat{i} \cdot (-2\hat{i} + \hat{j} + \hat{k}) = \lambda(7\hat{i} + 8\hat{j}) \cdot (-2\hat{i} + \hat{j} + \hat{k})$$

$$= -\frac{1}{2}(-14 + 8) + 2 = 5$$

7. Consider three vectors  $\vec{a}, \vec{b}, \vec{c}$ . Let  $|\vec{a}| = 2, |\vec{b}| = 3$

and  $\vec{a} = \vec{b} \times \vec{c}$ . If  $\alpha \in \left[0, \frac{\pi}{3}\right]$  is the angle between

the vectors  $\vec{b}$  and  $\vec{c}$ , then the minimum value of


$27|\vec{c} - \vec{a}|^2$  is equal to :

- (1) 110 (2) 105  
 (3) 124 (4) 121

Ans. (3)

**Sol.**  $|\vec{c} - \vec{a}| = |\vec{c}|^2 + |\vec{a}|^2 - 2\vec{a} \cdot \vec{c}$   
 $= |\vec{c}|^2 + 4 - 0$   
 $\therefore \vec{a} = \vec{b} \times \vec{c}$   
 $|\vec{a}| = |\vec{b} \times \vec{c}|$   
 $2 = 3|\vec{c}|\sin\alpha$   
 $|\vec{c}| = \frac{2}{3} \operatorname{cosec}\alpha \quad \alpha \in \left[0, \frac{\pi}{3}\right]$   
 $|\vec{c}|_{\min} = \frac{2}{3} \times \frac{2}{\sqrt{3}} \quad \operatorname{cosec}\alpha \in \left[\frac{2}{\sqrt{3}}, \infty\right)$   
 $\Rightarrow 27|\vec{c} - \vec{a}|_{\min}^2 = 27\left(\frac{16}{27} + 4\right) = 124$

- 8.** Let A(-1, 1) and B(2, 3) be two points and P be a variable point above the line AB such that the area of  $\Delta PAB$  is 10. If the locus of P is  $ax + by = 15$ , then  $5a + 2b$  is :
- (1)  $-\frac{12}{5}$  (2)  $-\frac{6}{5}$   
 (3) 4 (4) 6



**Sol.**

$$\frac{1}{2} \begin{vmatrix} h & k & 1 \\ 1 & 1 & 1 \\ 2 & 3 & 1 \end{vmatrix} = 10$$


$$-2x + 3y = 25$$

$$-\frac{6}{5}x + -y = 15$$

$$a = -\frac{6}{5}, b = \frac{9}{5}$$

$$5a = -6, 2b = \frac{18}{5}$$

- 9.** Let  $(\alpha, \beta, \gamma)$  be the point (8, 5, 7) in the line  $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-2}{5}$ . Then  $\alpha + \beta + \gamma$  is equal to
- (1) 16 (2) 18  
 (3) 14 (4) 20



$$\overline{AM} \cdot (2\hat{i} + 3\hat{j} + 5\hat{k}) = 0$$

$$(2\lambda - 7)(2) + (3\lambda - 6)(3) + (5\lambda - 5)(5) = 0$$

$$38\lambda = 57$$

$$\lambda = \frac{3}{2}$$

$$M\left(4, \frac{7}{2}, \frac{19}{2}\right)$$

$$A'(0, 2, 12)$$

- 10.** If the constant term in the expansion of  $\left(\frac{\sqrt[5]{3}}{x} + \frac{2x}{\sqrt[3]{5}}\right)^{12}$ ,  $x \neq 0$ , is  $\alpha \times 2^8 \times \sqrt[5]{3}$ , then  $25\alpha$  is equal to :
- (1) 639 (2) 724  
 (3) 693 (4) 742
- Ans. (3)**

**Sol.**

$$T_{r+1} = {}^{12}C_r \left(\frac{3^{1/5}}{x}\right)^{12-r} \left(\frac{2x}{5^{1/3}}\right)^r$$

$$T_{r+1} = \frac{{}^{12}C_r (3)^{\frac{12-r}{5}} (2)^r (x)^{2r-12}}{(5)^{r/3}}$$

$$r = 6$$

$$T_7 = \frac{{}^{12}C_6 (3)^{6/5} (2)^6}{5^2} \left(\frac{9 \times 11 \times 7}{25}\right) 2^8 \cdot 3^{1/5}$$

$$25\alpha = 693$$


11. Let  $f, g : \mathbb{R} \rightarrow \mathbb{R}$  be defined as :  $f(x) = |x - 1|$  and  $g(x) = \begin{cases} e^x, & x \geq 0 \\ x + 1, & x < 0 \end{cases}$ . Then the function  $f(g(x))$  is
- (1) neither one-one nor onto.
  - (2) one-one but not onto.
  - (3) both one-one and onto.
  - (4) onto but not one-one.

**Ans. (1)**

**Sol.**  $f(g(x)) = |g(x) - 1|$

$$f \circ g \begin{cases} |e^x - 1| & x \geq 0 \\ |x + 1 - 1| & x < 0 \end{cases}$$

$$f \circ g \begin{cases} e^x - 1 & x \geq 0 \\ -x & x < 0 \end{cases}$$



12. Let the circle  $C_1 : x^2 + y^2 - 2(x + y) + 1 = 0$  and  $C_2$  be a circle having centre at  $(-1, 0)$  and radius 2. If the line of the common chord of  $C_1$  and  $C_2$  intersects the y-axis at the point P, then the square of the distance of P from the centre of  $C_1$  is :

- (1) 2
- (2) 1
- (3) 6
- (4) 4

**Ans. (1)**

**Sol.**  $S_1 : x^2 + y^2 - 2x - 2y + 1 = 0$

$S_2 : x^2 + y^2 + 2x - 3 = 0$

Common chord =  $S_1 - S_2 = 0$

$-4x - 2y + 4 = 0$

$2x + y = 2 \Rightarrow P(0, 2)$


$d_{(c,p)}^2 = (1 - 0)^2 + (2 - 1)^2 = 2$

13. Let the set  $S = \{2, 4, 8, 16, \dots, 512\}$  be partitioned into 3 sets A, B, C with equal number of elements such that  $A \cup B \cup C = S$  and  $A \cap B = B \cap C = A \cap C = \phi$ . The maximum number of such possible partitions of S is equal to :

- (1) 1680
- (2) 1520
- (3) 1710
- (4) 1640

**Ans. (1)**

**Sol.**



$$\frac{9!}{(3!3!3!)} \times 3!$$

14. The values of m, n, for which the system of equations

$$\begin{aligned} x + y + z &= 4, \\ 2x + 5y + 5z &= 17, \\ x + 2y + mz &= n \end{aligned}$$

has infinitely many solutions, satisfy the equation :

- (1)  $m^2 + n^2 - m - n = 46$
- (2)  $m^2 + n^2 + m + n = 64$
- (3)  $m^2 + n^2 + mn = 68$
- (4)  $m^2 + n^2 - mn = 39$

**Ans. (4)**

**Sol.**  $D = \begin{vmatrix} 1 & 1 & 1 \\ 2 & 5 & 5 \\ 1 & 2 & m \end{vmatrix} = 0 \Rightarrow m = 2$

$D_3 = \begin{vmatrix} 1 & 1 & 4 \\ 2 & 5 & 17 \\ 1 & 2 & n \end{vmatrix} = 0 \Rightarrow n = 7$

15. The coefficients a, b, c in the quadratic equation  $ax^2 + bx + c = 0$  are from the set  $\{1, 2, 3, 4, 5, 6\}$ . If the probability of this equation having one real root bigger than the other is p, then 216p equals :

- (1) 57
- (2) 38
- (3) 19
- (4) 76

**Ans. (2)**



19. If  $y(\theta) = \frac{2\cos\theta + \cos 2\theta}{\cos 3\theta + 4\cos 2\theta + 5\cos\theta + 2}$ ,  
then at  $\theta = \frac{\pi}{2}$ ,  $y'' + y' + y$  is equal to:

- (1)  $\frac{3}{2}$  (2) 1  
(3)  $\frac{1}{2}$  (4) 2

Ans. (4)

Sol.  $y = \frac{2\cos\theta + 2\cos^2\theta - 1}{4\cos^3\theta - 3\cos\theta + 8\cos^2\theta - 4 + 5\cos\theta + 2}$

$$y = \frac{(2\cos^2\theta + 2\cos\theta - 1)}{(2\cos^2\theta + 2\cos\theta - 1)(2\cos\theta + 2)}$$

$$y = \frac{1}{2(1 + \cos\theta)}$$

$$\Rightarrow \theta = \frac{\pi}{2} \quad y = \frac{1}{2}$$

$$y' = \frac{1}{2} \left( \frac{-1}{(1 + \cos\theta)^2} \right) \times (-\sin\theta)$$

$$\Rightarrow \theta = \frac{\pi}{2} \quad y' = \frac{1}{2}$$

$$y'' = \frac{1}{2} \left[ \frac{\cos\theta(1 + \cos\theta)^2 - \sin\theta(2)(1 + \cos\theta)(-\sin\theta)}{(1 + \cos\theta)^4} \right]$$

$$\Rightarrow \theta = \frac{\pi}{2} \quad y'' = 1$$

20. For  $x \geq 0$ , the least value of K, for which  $4^{1+x} + 4^{1-x}$ ,  $\frac{K}{2}$ ,  $16^x + 16^{-x}$  are three consecutive terms of an A.P. is equal to:

- (1) 10 (2) 4  
(3) 8 (4) 16

Ans. (1)

Sol.  $k = 4 \left( 4^x + \frac{1}{4^x} \right) + \left( 4^{2x} + \frac{1}{4^{2x}} \right)$   
 $\geq 2$

$$k \geq 10$$

SECTION-B

21. Let the mean and the standard deviation of the probability distribution

X	$\alpha$	1	0	-3
P(X)	$\frac{1}{3}$	K	$\frac{1}{6}$	$\frac{1}{4}$

be  $\mu$  and  $\sigma$ , respectively. If  $\sigma - \mu = 2$ , then  $\sigma + \mu$  is equal to \_\_\_\_\_.

Ans. (5)

Sol.  $\frac{1}{3} + k + \frac{1}{6} + \frac{1}{4} = 1 \Rightarrow k = \frac{1}{4}$

$$\mu = \frac{\alpha}{3} + \frac{1}{4} - \frac{3}{4}$$

$$\mu = \frac{\alpha}{3} - \frac{1}{4}$$

$$\sigma = \sqrt{\left( \alpha^2 \frac{1}{3} + \frac{1}{4} + 9 \frac{1}{4} \right) - \left( \frac{\alpha}{3} - \frac{1}{4} \right)^2}$$

$$\sigma = \sqrt{\frac{2\alpha^2}{9} + \frac{1}{3} + \frac{1}{4}}$$

$$\sigma = \mu + 2$$

$$\sigma^2 = (\mu + 2)^2 \Rightarrow \frac{2\alpha^2}{9} + \frac{\alpha}{3} + \frac{9}{4} = \frac{\alpha}{9} + \frac{9}{4} + \alpha$$

$$\frac{\alpha^2}{9} - \frac{2}{3} = 0$$

$$\alpha = 0, \text{ (reject) or } \alpha = 6$$

( $\because x = 0$  is already given)

$$\Rightarrow \sigma + \mu = 2\mu + 2$$

$$= 5$$

22. Let  $y = y(x)$  be the solution of the differential

equation  $\frac{dy}{dx} + \frac{2x}{(1+x^2)^2} y = x e^{\frac{1}{1+x^2}}$ ;  $y(0) = 0$ .

Then the area enclosed by the curve

$f(x) = y(x) e^{-\frac{1}{1+x^2}}$  and the line  $y - x = 4$  is \_\_\_\_\_.

Ans. (18)

Sol.  $IF = e^{\int \frac{2x}{(1+x^2)^2} dx} = \frac{-}{1+x^2}$


$$y \cdot e^{\frac{-1}{1+x^2}} = \int x \cdot e^{\frac{1}{1+x^2}} \cdot e^{\frac{-1}{1+x^2}} dx$$

$$y \cdot e^{\frac{-1}{1+x^2}} = \frac{x^2}{2} + c$$

$$(0, 0) \Rightarrow \boxed{C=0}$$

$$y(x) = \frac{x^2}{2} e^{\frac{1}{1+x^2}}$$

$$f(x) = \frac{x^2}{2}$$




$$A = \int_{-2}^4 (x+4) - \frac{x^2}{2} dx = 18$$

23. The number of solutions of  $\sin^2 x + (2 + 2x - x^2)\sin x - 3(x - 1)^2 = 0$ , where  $-\pi \leq x \leq \pi$ , is


Ans. (2)

Sol.  $\sin^2 x - (x^2 - 2x - 2)\sin x - 3(x - 1)^2 = 0$   
 $\sin^2 x - (x - 1)^2 \sin x - 3(x - 1)^2 = 0$



$\sin x = -3$  (reject) or  $(x - 1)^2$

$\sin x = (x - 1)^2$




24. Let the point  $(-1, \alpha, \beta)$  lie on the line of the shortest distance between the lines  $\frac{x+2}{-3} = \frac{y-2}{4} = \frac{z-5}{2}$  and  $\frac{x+2}{-1} = \frac{y+6}{2} = \frac{z-1}{0}$ .

Then  $(\alpha - \beta)^2$  is equal to \_\_\_\_\_.

Ans. (25)

Sol.



$P(-3\lambda - 2, 4\lambda + 2, 2\lambda + 5)$

$Q(-\mu - 2, 2\mu - 6, 1)$

DRS of PQ =  $(3\lambda - \mu, 2\mu - 4\lambda - 8, -2\lambda - 4)$

$$\text{DRS of PQ} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 2 & 0 \\ -3 & 4 & 2 \end{vmatrix}$$

$= (4\hat{i} + 2\hat{j} + 2\hat{k})$

OR

$(2, 1, 1)$

$$\frac{3\lambda - \mu}{2} = \frac{2\mu - 4\lambda - 8}{1} = \frac{-2\lambda - 4}{1}$$

$\Rightarrow \mu = \lambda + 2$  &  $7\lambda = \mu - 8$

$\boxed{\lambda = -1} \quad \boxed{\mu = 1}$

$Q : (-3, -4, 1)$

$$L_{PQ} = \frac{x+3}{2} = \frac{y+4}{1} = \frac{z-1}{1}$$

$(-1, \alpha, \beta) \Rightarrow 1 = \frac{\alpha+4}{1} = \frac{\beta-1}{1}$

$\Rightarrow \alpha = -3, \beta = 2$

$(\alpha - \beta)^2 = 25$

25. If  $1 + \frac{\sqrt{3}-\sqrt{2}}{2\sqrt{3}} + \frac{5-2\sqrt{6}}{18} + \frac{9\sqrt{3}-11\sqrt{2}}{36\sqrt{3}} + \frac{49-20\sqrt{6}}{180} + \dots$   
upto  $\infty = 2 \sqrt{\frac{b}{a} + 1} \log_e \left( \frac{a}{b} \right)$ , where a and b are integers with  $\gcd(a, b) = 1$ , then  $11a + 18b$  is equal to \_\_\_\_\_.

Ans. (76)

Sol.  $S = 1 + \frac{x}{2\sqrt{3}} + \frac{x^2}{18} + \frac{x^3}{36\sqrt{3}} + \frac{x^4}{180} + \dots \infty$

Put  $\frac{x}{\sqrt{3}} = t$ , where  $x = \sqrt{3} - \sqrt{2}$

$$S = 1 + \frac{t}{2} + \frac{t^2}{6} + \frac{t^3}{12} + \frac{t^4}{20} + \dots$$

$$S = 1 + t \left( 1 - \frac{1}{2} \right) + t^2 \left( \frac{1}{2} - \frac{1}{3} \right) + t^3 \left( \frac{1}{3} - \frac{1}{4} \right) + t^4 \left( \frac{1}{4} - \frac{1}{5} \right) + \dots$$

$$S = \left( 1 + t + \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{4} + \dots \right) - \left( \frac{t}{2} + \frac{t^2}{3} + \frac{t^3}{4} + \frac{t^4}{5} + \dots \right)$$

$$S = \left( t + \frac{t^2}{2} + \dots \right) - \frac{1}{t} \left( t + \frac{t^2}{2} + \frac{t^3}{3} + \dots \right) + 2$$

$$S = 2 + \left( 1 - \frac{1}{t} \right) (-\log(1-t)) = \left( -1 \right) \log(1-t) + 2$$

$$S = 2 \left( \frac{\sqrt{3}}{\sqrt{3}-\sqrt{2}} - 1 \right) \log \left( 1 - \frac{\sqrt{3}-\sqrt{2}}{\sqrt{3}} \right)$$

$$S = 2 \left( \frac{\sqrt{2}}{\sqrt{3}-\sqrt{2}} \log_e \frac{\sqrt{3}}{\sqrt{3}-\sqrt{2}} \right)$$

$$S = 2 + \frac{(\sqrt{6}+2)}{2} \log_e \frac{2}{3} = 2 + \left( \frac{\sqrt{3}}{2} + 1 \right) \log_e \frac{2}{3}$$

$a = 2, b = 3$

$11a + 18b = 11 \times 2 + 18 \times 3 = 76$

26. Let  $a > 0$  be a root of the equation  $2x^2 + x - 2 = 0$ .

If  $\lim_{x \rightarrow \frac{1}{a}} \frac{16(1 - \cos(2 + x - 2x^2))}{(1 - ax^2)} = \alpha + \beta\sqrt{17}$ , where

$\alpha, \beta \in \mathbb{Z}$  then  $\alpha + \beta$  is equal to \_\_\_\_\_.

Ans. (170)

Sol.  $2x^2 + x - 2 = 0 \begin{cases} a \\ b \end{cases}$

$$2x^2 - x - 2 = 0 \begin{cases} \frac{1}{a} \\ \frac{1}{b} \end{cases}$$

$$\lim_{x \rightarrow \frac{1}{a}} 16 \cdot \frac{\left( 1 - \cos 2 \left( x - \frac{1}{a} \right) \left( x - \frac{1}{b} \right) \right)}{4 \left( x - \frac{1}{b} \right)^2} \times \frac{4 \left( x - \frac{1}{b} \right)^2}{a^2 \left( x - \frac{1}{a} \right)^2}$$

$$= 16 \times \frac{2}{a^2} \left( \frac{1}{a} - \frac{1}{b} \right)^2 = \frac{32}{a^2} \left( \frac{17}{4} \right) = \frac{17.8}{2} = \frac{17 \times 8 \times 16}{(-1 + \sqrt{117})^2}$$

$$= \frac{136.16}{18.2\sqrt{7}} \frac{18 + 2\sqrt{7}}{18 + 2\sqrt{7}}$$

$$= \frac{136}{256} (18 + 2\sqrt{7}) \cdot 16$$

$$= 153 + 17\sqrt{17} = \alpha + \beta\sqrt{17}$$

$\alpha + \beta = 153 + 17 = 170$

27. If  $f(t) = \int_0^{\pi} \frac{2x dx}{1 - \cos t \sin^2 x}$ ,  $0 < t < \pi$ , then the value

of  $\int_0^{\frac{\pi}{2}} \frac{\pi^2 dt}{f(t)}$  equals \_\_\_\_\_.

Ans. (1)

Sol.  $f(t) = \int_0^{\pi} \frac{2x}{1 - \cos t \sin^2 x} dx \dots (1)$



$$= 2 \int_0^{\pi} \frac{(\pi - x) dx}{1 - \cos^2 t \sin^2 x} \quad \dots(2)$$

$$2f(t) = 2 \int_0^{\pi} \frac{\pi}{1 - \cos^2 t \sin^2 x} dx$$

$$f(t) = \int_0^{\pi} \frac{\pi}{1 - \cos^2 t \sin^2 x} dx$$

divide & by  $\cos^2 x$

$$f(t) = \pi \int_0^{\pi} \frac{\sec^2 x dx}{\sec^2 x - \cos^2 t \tan^2 x}$$

$$f(t) = 2 \int_0^{\pi/2} \frac{\sec^2 x dx}{\sec^2 x - \cos^2 t \tan^2 x}$$

$$\tan x = z$$

$$\sec^2 x dx = dz$$

$$f(t) = 2 \int_0^{\infty} \frac{dz}{1 + \sin^2 t \cdot z^2}$$

$$= \frac{\pi^2}{\sin t}$$

$$\text{Then } \int_0^{\pi/2} \frac{\pi^2}{f(t)} dt$$

$$= \int_0^{\pi/2} \sin t dt$$

$$= 1$$

28. Let the maximum and minimum values of  $(\sqrt{8x - x^2 - 12} - 4)^2 + (x - 7)^2$ ,  $x \in \mathbb{R}$  be  $M$  and  $m$  respectively. Then  $M^2 - m^2$  is equal to \_\_\_\_\_.


Ans. (1600)

Sol.  $(x - 7)^2 + (y - 4)^2$

$$y = \sqrt{8x - x^2 - 12}$$

$$y = -(x - 4)^2 + 16 - 12$$

$$(x - 4)^2 + y^2 = 4$$



$$m = 9$$

$$M = 41$$

$$M^2 - m^2 = 41^2 - 9^2 = 1600$$

29. Let a line perpendicular to the line  $2x - y = 10$  touch the parabola  $y^2 = 4(x - 9)$  at the point  $P$ . The distance of the point  $P$  from the centre of the circle  $x^2 + y^2 - 14x - 8y + 56 = 0$  is \_\_\_\_\_.

Ans. (10)

Sol.  $y^2 = 4(x - 9)$

$$\text{slope of tangent} = \frac{-}{2}$$

$$\text{Point of contact } P \left( 9 + \frac{1}{\left(-\frac{1}{2}\right)^2}, \frac{2 \times 1}{\frac{1}{2}} \right)$$

$$P(13, -4)$$

$$\text{center of circle } C(7, 4)$$

$$\text{distance } CP = \sqrt{(13 - 7)^2 + (-4 - 4)^2} = 10$$

30. The number of real solutions of the equation  $x|x + 5| + 2|x + 7| - 2 = 0$  is \_\_\_\_\_.

Ans. (3)

30. The number of real solutions of the equation  $x|x + 5| + 2|x + 7| - 2 = 0$  is \_\_\_\_\_.

Ans. (3)

Sol. Case I :  $x \geq -5$

$$x^2 + 5x + 2x + 12 = 0$$

$$x^2 + 7x + 12 = 0$$

$$x = -3, -4$$

Case II :  $-7 < x < -5$

$$-x^2 - 5x + 2x + 14 - 2 = 0$$



$$-x^2 - 3x + 12 = 0$$

$$x = \frac{-3 \pm \sqrt{9 + 48}}{2}$$

$$= \frac{-3 \pm \sqrt{57}}{2}$$

$$x = \frac{-3 - \sqrt{57}}{2}, \frac{-3 + \sqrt{57}}{2} \text{ (rejected)}$$

**Case III :**  $x \leq -7$

$$-x^2 - 5x - 2x - 14 - 2 = 0$$

$$x^2 + 7x + 16 = 0$$

$$D = 49 - 64 < 0$$

No solutions

No. of solutions = 3







36. If  $n$  is the number density and  $d$  is the diameter of the molecule, then the average distance covered by a molecule between two successive collisions (i.e. mean free path) is represented by :

$$(1) \frac{1}{\sqrt{2n\pi d^2}} \quad (2) \sqrt{2n\pi d^2}$$

$$(3) \frac{1}{\sqrt{2n\pi d^2}} \quad (4) \frac{1}{\sqrt{2n^2\pi^2 d^2}}$$

Ans. (3)

Sol.  $n$  = number of molecule per unit volume  
 $d$  = diameter of the molecule

$$\lambda = \frac{1}{\sqrt{2n\pi d^2 n}} \quad (\text{By Theory})$$

37. A particle moves in x-y plane under the influence of a force  $\vec{F}$  such that its linear momentum is  $\vec{P}(t) = \hat{i} \cos(kt) - \hat{j} \sin(kt)$ . If  $k$  is constant, the angle between  $\vec{F}$  and  $\vec{P}$  will be :

$$(1) \frac{\pi}{2} \quad (2) \frac{\pi}{6}$$

$$(3) \frac{\pi}{4} \quad (4) \frac{\pi}{3}$$

Ans. (1)

Sol.  $\vec{P} = \cos(kt)\hat{i} - \sin(kt)\hat{j}$  ;  $|\vec{P}| =$

$$\therefore \vec{P} = m\vec{v}$$

$$\therefore \hat{P} = \hat{v}$$

$$\Rightarrow \hat{v} = \cos(kt)\hat{i} - \sin(kt)\hat{j}$$

$$\hat{a} = \frac{-k \sin(kt)\hat{i} - k \cos(kt)\hat{j}}{k}$$

$$\Rightarrow \hat{a} = -\sin kt \hat{i} - \cos kt \hat{j}$$

$$\therefore \hat{F} = \hat{a} = -\sin kt \hat{i} - \cos kt \hat{j}$$

$$\cos \theta = \frac{\hat{F} \cdot \hat{P}}{|\hat{F}| |\hat{P}|} = -\frac{\sin kt \cos t + \sin kt \cos t}{1 \times 1} = 0$$

$$\Rightarrow \theta = \frac{\pi}{2}$$

38. The electrostatic force ( $\vec{F}_1$ ) and magnetic force ( $\vec{F}_2$ ) acting on a charge  $q$  moving with velocity  $\vec{v}$  can be written :

$$(1) \vec{F}_1 = q\vec{V} \cdot \vec{E}, \vec{F}_2 = q(\vec{B} \cdot \vec{V})$$

$$(2) \vec{F}_1 = q\vec{B}, \vec{F}_2 = q(\vec{B} \times \vec{V})$$

$$(3) \vec{F}_1 = q\vec{E}, \vec{F}_2 = q(\vec{V} \times \vec{B})$$

$$(4) \vec{F}_1 = q\vec{E}, \vec{F}_2 = q(\vec{B} \times \vec{V})$$

Ans. (3)

Sol.  $\vec{F}_1 = q\vec{E}$  (Theory)

$$\vec{F}_2 = q(\vec{V} \times \vec{B})$$

39. A man carrying a monkey on his shoulder does cycling smoothly on a circular track of radius 9m and completes 120 revolutions in 3 minutes. The magnitude of centripetal acceleration of monkey is (in  $\text{m/s}^2$ ) :

$$(1) \text{zero} \quad (2) 16 \pi^2 \text{ms}^{-2}$$

$$(3) 4\pi^2 \text{ms}^{-2} \quad (4) 57600 \pi^2 \text{ms}^{-2}$$

Ans. (2)

Sol. Given :  $R = 9\text{m}$ ,

120 revolution in 3 min

$$\omega = \frac{120 \text{ Rev.}}{3 \text{ min.}} = \frac{120 \times 2\pi \text{ rad}}{3 \times 60 \text{ sec}} = \frac{4\pi}{3} \text{ rad/s}$$

$$a_{\text{centripetal}} = \omega^2 R = \left(\frac{4\pi}{3}\right)^2 \times 9 = 16\pi^2 \text{m/s}^2$$

40. A series LCR circuit is subjected to an AC signal of 200 V, 50 Hz. If the voltage across the inductor ( $L = 10 \text{ mH}$ ) is 31.4 V, then the current in this circuit is \_\_\_\_\_ :

$$(1) 68 \text{ A} \quad (2) 63 \text{ A}$$

$$(3) 10 \text{ A} \quad (4) 10 \text{ mA}$$

Ans. (3)

Sol. Voltage across inductor  $V_L = IX_L$

$$31.4 = I[L\omega]$$

$$31.4 = I[L(2\pi f)]$$

$$31.4 = I[10 \times 10^{-3}(2 \times 3.14) \times 50]$$

$$\Rightarrow I = 10 \text{ A}$$

41. What is the dimensional formula of  $ab^{-1}$  in the equation  $\left(P + \frac{a}{V^2}\right)(V - b) = RT$ , where letters have their usual meaning.

- (1)  $[M^0L^3T^{-2}]$                       (2)  $[ML^2T^{-2}]$   
 (3)  $[M^{-1}L^5T^3]$                       (4)  $[M^6L^7T^4]$

Ans. (2)


Sol.  $\therefore [V] = [b]$   
 $\therefore$  Dimension of  $b = [L^3]$

&  $[P] = \left[\frac{a}{V^2}\right]$

$[a] = [PV^2] = [ML^{-1}T^{-2}][L^6]$   
 Dimension of  $a = [ML^5T^{-2}]$

$\therefore ab^{-1} = \frac{[ML^5T^{-2}]}{[L^3]} = [ML^2T^{-2}]$


42. The output (Y) of logic circuit given below is 0 only when :



- (1)  $A = 1, B = 0$                       (2)  $A = 0, B = 0$   
 (3)  $A = 1, B = 1$                       (4)  $A = 0, B = 1$

Ans. (2)

Sol.



43. A body is moving unidirectionally under the influence of a constant power source. Its displacement in time  $t$  is proportional to :

- (1)  $t^2$                                       (2)  $t^{2/3}$   
 (3)  $t^{3/2}$                                   (4)  $t$

Ans. (3)

Sol.  $P = \text{constant} \Rightarrow FV = \text{constant}$

$\Rightarrow m \frac{dV}{dt} = \text{constant}$

$\int_0^V V dV = (C) \int dt$

$\left(\frac{V^2}{2}\right) = Ct$

$V \propto t^{1/2}$

$\frac{ds}{dt} \propto t^{1/2}$

$\int_0^s ds = K \int t^{1/2} dt$

$S = K \times \frac{2}{3} t^{3/2}$

$S \propto t^{3/2}$

$\therefore$  displacement is proportional to  $(t)^{3/2}$

44. Match List-I with List-II :-

	List-I EM-Wave		List-II Wavelength Range
(A)	Infra-red	(I)	$< 10^{-3}$ nm
(B)	Ultraviolet	(II)	400 nm to 1 nm
(C)	X-rays	(III)	1 mm to 700 nm
(D)	Gamma rays	(IV)	1 nm to $10^{-3}$ nm

Choose the correct answer from the options given below :

- (1) (A)-(II), (B)-(I), (C)-(IV), (D)-(III)  
 (2) (A)-(III), (B)-(II), (C)-(IV), (D)-(I)  
 (3) (A)-(IV), (B)-(III), (C)-(II), (D)-(I)  
 (4) (A)-(I), (B)-(III), (C)-(II), (D)-(IV)

Ans. (2)

Sol. Infrared is the least energetic thus having biggest wavelength ( $\lambda$ ) & gamma rays are most energetic thus having smallest wavelength ( $\lambda$ ).




48. A heavy box of mass 50 kg is moving on a horizontal surface. If coefficient of kinetic friction between the box and horizontal surface is 0.3 then force of kinetic friction is :

- (1) 14.7 N
- (2) 147 N
- (3) 1.47 N
- (4) 1470 N

Ans. (2)

Sol.



$$F_k = \mu_k N = 0.3 \times 50 \times 9.8 = 147 \text{ N}$$

49. A satellite revolving around a planet in stationary orbit has time period 6 hours. The mass of planet is one-fourth the mass of earth. The radius orbit of planet is : (Given = Radius of geo-stationary orbit for earth is  $4.2 \times 10^4 \text{ km}$ )

- (1)  $1.4 \times 10^4 \text{ km}$
- (2)  $8.4 \times 10^4 \text{ km}$
- (3)  $1.68 \times 10^5 \text{ km}$
- (4)  $1.05 \times 10^4 \text{ km}$

Ans. (4)


Sol.  $T = \frac{2\pi r^{3/2}}{\sqrt{GM}}$

$$\frac{T_1}{T_2} = \left(\frac{r_1}{r_2}\right)^{3/2} \left(\frac{M_2}{M_1}\right)^{1/2}$$

$$\frac{6}{24} = \left(\frac{r_1}{4.2 \times 10^4}\right)^{3/2} \left(\frac{M}{M/4}\right)^{1/2}$$

$$r_1 = 1.05 \times 10^4 \text{ km}$$


50. The ratio of heat dissipated per second through the resistance  $5 \Omega$  and  $10 \Omega$  in the circuit given below is :



- (1) 1 : 2
- (2) 2 : 1
- (3) 4 : 1
- (4) 1 : 1

Ans. (2)

Sol.



$$\frac{i_1}{i_2} = \frac{10}{5} = \frac{2}{1}$$

$$\frac{P_1}{P_2} = \frac{i_1^2 R_1}{i_2^2 R} = \left(\frac{2}{1}\right)^2 \times \frac{5}{10} = \frac{2}{1}$$

**SECTION-B**

51. A solenoid of length 0.5 m has a radius of 1 cm and is made up of 'm' number of turns. It carries a current of 5A. If the magnitude of the magnetic field inside the solenoid is  $6.28 \times 10^{-3} \text{ T}$ , then the value of m is :

Ans. (500)

Sol.  $\mu_0 n i = B$  n = number of turns per unit length

$$\mu_0 \left(\frac{m}{l}\right) i = B$$


$$m = \frac{B \cdot l}{\mu_0 i} = \frac{6.28 \times 10^{-3} \cdot 0.5}{12.56 \cdot 10^{-7} \times 5}$$

$$m = 500$$

52. The shortest wavelength of the spectral lines in the Lyman series of hydrogen spectrum is 915 Å. The longest wavelength of spectral lines in the Balmer series will be \_\_\_\_\_ Å.

Ans. (6588)

Sol. Lyman Series



$$\text{Shortest, } \frac{hc}{\lambda} = -13.6 \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$\lambda \downarrow E \uparrow ; \frac{hc}{\lambda_0} = -13.6(1)$$

Balmer Series :

$$\text{_____ } n = 3$$

$$\text{_____ } n = 2$$

$$\frac{hc}{\lambda_1} = -13.6 \left( \frac{1}{2^2} - \frac{1}{3^2} \right)$$

$$\frac{hc}{\lambda_1} = -13.6 \left( \frac{1}{4} - \frac{1}{9} \right)$$

$$\frac{hc}{\lambda_1} = -13.6 \times \left( \frac{5}{36} \right)$$


$$\Rightarrow \frac{-13.6\lambda_0}{\lambda_1} = -13.6 \times \frac{5}{36}$$

$$\lambda_1 = \frac{\lambda_0 \times 36}{5} = \frac{915 \times 36}{5} = 6588$$

53. In a single slit experiment, a parallel beam of green light of wavelength 550 nm passes through a slit of width 0.20 mm. The transmitted light is collected on a screen 100 cm away. The distance of first order minima from the central maximum will be  $x \times 10^{-5}$  m. The value of x is :

Ans. (275)

Sol.




$$y = \frac{\lambda D}{d} = \frac{550 \times 10^{-9} \times 100 \times 10^{-2}}{0.2 \times 10^{-3}} = 275$$

54. A sonometer wire of resonating length 90 cm has a fundamental frequency of 400 Hz when kept under some tension. The resonating length of the wire with fundamental frequency of 600 Hz under same tension \_\_\_\_\_ cm.

Ans. (60)

Sol.



$$f_0 = 400 \text{ Hz ; } v = \sqrt{\frac{T}{\mu}} = \text{constant}$$

$$\frac{\lambda}{2} = L ; v = f_0 \lambda$$

$$\frac{v}{2f_0} = L \Rightarrow v = 2Lf_0$$

$$L' = \frac{v}{2f'} = \frac{2Lf_0}{2f'}$$

$$= \frac{Lf_0}{f'} = \frac{90 \times 400}{600} = 60$$

55. A hollow sphere is rolling on a plane surface about its axis of symmetry. The ratio of rotational kinetic energy to its total kinetic energy is  $\frac{x}{5}$ . The value of x is \_\_\_\_\_.

Ans. (2)


$$\text{Sol. } \frac{\frac{1}{2} I \omega^2}{\frac{1}{2} I \omega^2 + \frac{1}{2} m v^2} = \frac{\left(\frac{1}{2}\right) \left(\frac{2}{3} m R^2\right) \omega^2}{\left(\frac{1}{2}\right) \left(\frac{2}{3} m R^2\right) \omega^2 + \frac{1}{2} m (R\omega)^2}$$

$$= \frac{\frac{2}{3}}{\frac{2}{3} + 1} = \frac{2}{5}$$

$$x = 2$$



56. A hydraulic press containing water has two arms with diameters as mentioned in the figure. A force of 10 N is applied on the surface of water in the thinner arm. The force required to be applied on the surface of water in the thicker arm to maintain equilibrium of water is \_\_\_\_\_ N.




Ans. (1000 N)

Sol. 
$$\frac{F_1}{A_1} = \frac{F_2}{A_2}$$

$$\frac{F_1}{\pi(7)^2} = \frac{10}{\pi \times (0.7)^2}$$

$$F_1 = 1000 \text{ N}$$

57. The electric field at point p due to an electric dipole is E. The electric field at point R on equatorial line will be  $\frac{E}{x}$ . The value of x :



Ans. (16)

Sol. 
$$E_p = \frac{2Kq}{r^3} = E$$

$$E_R = \frac{Kq}{(2r)^3} = \frac{E}{16}$$

$$x = 16$$

58. The maximum height reached by a projectile is 64 m. If the initial velocity is halved, the new maximum height of the projectile is \_\_\_\_\_ m.

Ans. (16)

Sol. 
$$H_{\max} = \frac{u^2 \sin^2 \theta}{2g}$$

$$\frac{H_{1\max}}{H_{2\max}} = \frac{u_1^2}{u_2^2}$$

$$\frac{64}{H_{2\max}} = \frac{u^2}{(u/2)^2}$$

$$H_{2\max} = 16 \text{ m}$$

59. A wire of resistance  $20 \Omega$  is divided into 10 equal parts. A combination of two parts are connected in parallel and so on. Now resulting pairs of parallel combination are connected in series. The equivalent resistance of final combination is \_\_\_\_\_  $\Omega$ .

Ans. (5)

Sol.

$$\boxed{20\Omega} \Rightarrow 10 \text{ equal part}$$

Each part has resistance =  $2\Omega$

2 parts are connected in parallel so,  $R = 1\Omega$

Now, there will be 5 parts each of resistance  $1\Omega$ , they are connected in series.

$$R_{\text{eq}} = 5R, R_{\text{eq}} = 5\Omega$$

60. The current in an inductor is given by  $I = (3t + 8)$  where t is in second. The magnitude of induced emf produced in the inductor is 12 mV. The self-inductance of the inductor \_\_\_\_\_ mH.

Ans. (4)

Sol.  $I = 3t + 8$

$$\varepsilon = 12 \text{ mV}$$











$$|\varepsilon| = L \left| \frac{dI}{dt} \right|$$


$$12 = L \times 3$$

$$L = 4 \text{ mH}$$

(Held On Friday 05<sup>th</sup> April, 2024)

TIME : 3 : 00 PM to 6 : 00 PM

CHEMISTRY	TEST PAPER WITH SOLUTION		
<b>SECTION-A</b>			
<p><b>61. Match List - I with List - II.</b></p> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top;"> <b>List - I</b>                      (A) ICl                      (B) ICl<sub>3</sub>                      (C) ClF<sub>5</sub>                      (D) IF<sub>7</sub> </td> <td style="width: 50%; vertical-align: top;"> <b>List - II</b>                      (I) T -Shape                      (II) Square pyramidal                      (III) Pentagonal bipyramidal                      (IV) Linear                 </td> </tr> </table> <p>Choose the <b>correct</b> answer from the options given below:</p> <p>(1) (A)–(I), (B)–(IV), C–(III), D–(II)                      (2) (A)–(I), (B)–(III), C–(II), D–(IV)                      (3) (A)–(IV), (B)–(I), C–(II), D–(III)                      (4) (A)–(IV), (B)–(III), C–(II), D–(I)</p> <p><b>Ans. (3)</b></p> <p><b>Sol.</b> A. I – Cl (iv) linear</p> <p>B.  (I) T-shape</p> <p>C.  (II) Square pyramidal</p> <p>D.  (III) Pentagonal bipyramidal</p> <p><b>62. While preparing crystals of Mohr's salt, dil. H<sub>2</sub>SO<sub>4</sub> is added to a mixture of ferrous sulphate and ammonium sulphate, before dissolving this mixture in water, dil. H<sub>2</sub>SO<sub>4</sub> is added here to:</b></p> <p>(1) prevent the hydrolysis of ferrous sulphate                      (2) prevent the hydrolysis of ammonium sulphate                      (3) make the medium strongly acidic                      (4) increase the rate of formation of crystals</p> <p><b>Ans. (1)</b></p>	<b>List - I</b> (A) ICl (B) ICl <sub>3</sub> (C) ClF <sub>5</sub> (D) IF <sub>7</sub>	<b>List - II</b> (I) T -Shape (II) Square pyramidal (III) Pentagonal bipyramidal (IV) Linear	<p><b>Sol.</b> Fe<sup>+2</sup> ions undergoes hydrolysis, therefore while preparing aqueous solution of ferrous sulphate and ammonium sulphate in water dilute sulphuric acid is added to prevent hydrolysis of ferrous sulphate.</p> <p><b>63. Identify the major product in the following reaction.</b></p> <p></p> <p>(1)                       (2)                       (3)                       (4) </p> <p><b>Ans. (3)</b></p> <p><b>Sol.</b> </p> <p><b>64. The correct nomenclature for the following compound is:</b></p> <p></p> <p>(1) 2-carboxy-4-hydroxyhept-6-enal                      (2) 2-carboxy-4-hydroxyhept-7-enal                      (3) 2-formyl-4-hydroxyhept-6-enoic acid                      (4) 2-formyl-4-hydroxyhept-7-enoic acid</p> <p><b>Ans. (3)</b></p>
<b>List - I</b> (A) ICl (B) ICl <sub>3</sub> (C) ClF <sub>5</sub> (D) IF <sub>7</sub>	<b>List - II</b> (I) T -Shape (II) Square pyramidal (III) Pentagonal bipyramidal (IV) Linear		



2-formyl-4-hydroxyhept-6-enoic acid

65. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.


**Assertion (A)** :  $\text{NH}_3$  and  $\text{NF}_3$  molecule have pyramidal shape with a lone pair of electrons on nitrogen atom. The resultant dipole moment of  $\text{NH}_3$  is greater than that of  $\text{NF}_3$ .

**Reason (R)** : In  $\text{NH}_3$ , the orbital dipole due to lone pair is in the same direction as the resultant dipole moment of the N-H bonds. F is the most electronegative element.


In the light of the above statements, choose the **correct** answer from the options given below:

- (1) Both (A) and (R) are true and (R) is the correct explanation of (A)
- (2) (A) is false but (R) is true
- (3) (A) is true but (R) is false
- (4) Both (A) and (R) are true but (R) is NOT the correct explanation of (A)

Ans. (1)



Resultant dipole moment =  $0.80 \times 10^{-30}$  Cm



Resultant dipole moment =  $4.90 \times 10^{-30}$  cm

66. Given below are two statements:

**Statement I** : On passing  $\text{HCl}_{(g)}$  through a saturated solution of  $\text{BaCl}_2$ , at room temperature white turbidity appears.

**Statement II** : When  $\text{HCl}$  gas is passed through a saturated solution of  $\text{NaCl}$ , sodium chloride is precipitated due to common ion effect.

In the light of the above statements, choose the **most appropriate** answer from the options given below:

- (1) **Statement I** is correct but **Statement II** is incorrect
- (2) Both **Statement I** and **Statement II** are incorrect
- (3) **Statement I** is incorrect but **Statement II** is correct
- (4) Both **Statement I** and **Statement II** are correct

Ans. (1)

Sol.  $\text{BaCl}_2$ ,  $\text{NaCl}$  are soluble but on adding  $\text{HCl}_{(g)}$  to  $\text{BaCl}_2$ ,  $\text{NaCl}$  solutions, Sodium or Barium chlorides may precipitate out, as a consequence of the law of mass action.

67. The metal atom present in the complex  $\text{MABXL}$  (where A, B, X and L are unidentate ligands and M is metal) involves  $sp^3$  hybridization. The number of geometrical isomers exhibited by the complex is:

- (1) 4
- (2) 0
- (3) 2
- (4) 3

Ans. (2)


Sol. Tetrahedral complex does not show geometrical isomerism.

68. Match **List - I** with **List - II**.

<b>List - I</b> (Pair of Compounds)	<b>List - II</b> (Isomerism)
(A) n-propanol and Isopropanol	(I) Metamerism
(B) Methoxypropane and ethoxyethane	(II) Chain Isomerism
(C) Propanone and propanal	(III) Position Isomerism
(D) Neopentane and Isopentane	(IV) Functional Isomerism

- (1) (A)–(II), (B)–(I), (C)–(IV), (D)–(III)
- (2) (A)–(III), (B)–(I), (C)–(II), (D)–(IV)
- (3) (A)–(I), (B)–(III), (C)–(IV), (D)–(II)
- (4) (A)–(III), (B)–(I), (C)–(IV), (D)–(II)

Ans. (4)



**69.** The quantity of silver deposited when one coulomb charge is passed through  $\text{AgNO}_3$  solution:

- (1) 0.1 g atom of silver
- (2) 1 chemical equivalent of silver
- (3) 1 g of silver
- (4) 1 electrochemical equivalent of silver

**Ans. (4)**


**Sol.**  $W = ZIt$

$$W = ZQ$$


$$Q = \frac{W}{Z}$$

$$W = ZQ = (\text{electrochemical equivalent})$$

**70.** Which one of the following reactions is NOT possible?



**Ans. (2)**



**71.** Given below are two statements :

**Statement I :** The metallic radius of Na is  $1.86 \text{ \AA}$  and the ionic radius of  $\text{Na}^+$  is lesser than  $1.86 \text{ \AA}$ .

**Statement II :** Ions are always smaller in size than the corresponding elements.

In the light of the above statements, choose the **correct** answer from the options given below :


- (1) **Statement I** is correct but **Statement II** is false
- (2) Both **Statement I** and **Statement II** are true
- (3) Both **Statement I** and **Statement II** are false
- (4) **Statement I** is incorrect but **Statement II** is true

**Ans. (1)**

**Sol.**  $r_{\text{Na}} > r_{\text{Na}^+}$

So, Statement (I) is correct but size of anions are greater than size of neutral atoms.


So statement (II) is incorrect.




Consider the above reaction sequence and identify the major product P.

- (1) Methane
- (2) Methanal
- (3) Methoxymethane
- (4) Methanoic acid

**Ans. (1)**




**73.** Consider the given chemical reaction :



Product "A" is :

- (1) picric acid
- (2) oxalic acid
- (3) acetic acid
- (4) adipic acid

**Ans. (4)**



74. For the electro chemical cell  
 $M|M^{2+}||X|X^{2-}$   
 If  $E^0_{(M^{2+}/M)} = 0.46\text{ V}$  and  $E^0_{(X/X^{2-})} = 0.34\text{ V}$ .

Which of the following is **correct** ?

- (1)  $E_{\text{cell}} = -0.80\text{ V}$   
 (2)  $M + X \rightarrow M^2 + X^{2-}$  is a spontaneous reaction  
 (3)  $M^{2+} + X^{2-} \rightarrow M + X$  is a spontaneous reaction  
 (4)  $E_{\text{cell}} = 0.80\text{ V}$

Ans. (3)

Sol.  $M | M^{2+} || X / X^{2-}$

$$E_{\text{cell}}^0 = E^0_{M/M^{2+}} + E^0_{X/X^{2-}}$$

$$= -0.46 + 0.34 = -0.12\text{ V}$$

As  $E_{\text{cell}}^0$  is negative so anode becomes cathode and cathode become anode. Spontaneous reaction will be  
 $M^{2+} + X^{2-} \rightarrow M + X$


75. The number of moles of methane required to produce 11g  $\text{CO}_2(\text{g})$  after complete combustion is:

(Given molar mass of methane in  $\text{g mol}^{-1}$  : 16)

- (1) 0.75 (2) 0.25  
 (3) 0.35 (4) 0.5

Ans. (2)

Sol.  $\text{C}_n\text{H}_{2n+2} + \frac{3n+1}{2}\text{O}_2 \rightarrow n\text{CO}_2 + (n+1)\text{H}_2\text{O}$



4gm 11gm

0.25 mole 0.25 mole

0.25 mole  $\text{CH}_4$  gives 0.25 mole (or 11gm)  $\text{CO}_2$

76. The number of complexes from the following with no electrons in the  $t_2$  orbital is \_\_\_\_\_.

$\text{TiCl}_4$ ,  $[\text{MnO}_4]^-$ ,  $[\text{FeO}_4]^{2-}$ ,  $[\text{FeCl}_4]^-$ ,  $[\text{CoCl}_4]^{2-}$

- (1) 3 (2) 1  
 (3) 4 (4) 2

Ans. (1)

Sol.  $\text{TiCl}_4 \Rightarrow \text{Ti}^{+4} \quad e^0 t_2^0$

$\text{MnO}_4^- \Rightarrow \text{Mn}^{+7} \quad e^0 t_2^0$

$\text{FeO}_4^{2-} \Rightarrow \text{Fe}^{+6} \quad e^2 t_2^0$

$\text{FeCl}_4^{2-} \Rightarrow \text{Fe}^{+2} \quad e^3 t_2^3$


$\text{CoCl}_4^{2-} \Rightarrow \text{Co}^{+2} \quad e^4 t_2^3$

77. The number of ions from the following that have the ability to liberate hydrogen from a dilute acid is \_\_\_\_\_.  $\text{Ti}^{2+}$ ,  $\text{Cr}^{2+}$  and  $\text{V}^{2+}$






- (1) 0 (2) 2  
 (3) 3 (4) 1

Ans. (3)

Sol. The ions  $\text{Ti}^{2+}$ ,  $\text{V}^{2+}$ ,  $\text{Cr}^{2+}$  are strong reducing agents and will liberate hydrogen from a dilute acid, eg.




78. Identify A and B in the given chemical reaction sequence :-



Ans. (2)

Sol.



79. The correct statements from the following are :
- (A) The decreasing order of atomic radii of group 13 elements is  $Tl > In > Ga > Al > B$ .
- (B) Down the group 13 electronegativity decreases from top to bottom.
- (C) Al dissolves in dil. HCl and liberate  $H_2$  but conc.  $HNO_3$  renders Al passive by forming a protective oxide layer on the surface.
- (D) All elements of group 13 exhibits highly stable +1 oxidation state.
- (E) Hybridisation of Al in  $[Al(H_2O)_6]^{3+}$  ion is  $sp^3d^2$ .

Choose the **correct** answer from the options given below :

- (1) (C) and (E) only  
 (2) (A), (C) and (E) only  
 (3) (A), (B), (C) and (E) only  
 (4) (A) and (C) only

**Ans. (1)**

**Sol.** A. size order  $Tl > In > Al > Ga > B$

B. Electronegativity order  $B > Al < Ga < In < Tl$

D. B, Al are more stable in +3 oxidation state

So, only C, E statements are correct.

80. Coagulation of egg, on heating is because of :

- (1) Denaturation of protein occurs  
 (2) The secondary structure of protein remains unchanged  
 (3) Breaking of the peptide linkage in the primary structure of protein occurs  
 (4) Biological property of protein remains unchanged

**Ans. (1)**

**Sol.** Coagulation of egg give primary structure of protein, which is known as denaturation of protein

### SECTION-B

81. Combustion of 1 mole of benzene is expressed at  $C_6H_6(1) + \frac{15}{2}O_2(g) \rightarrow CO_2(g) + 3H_2O(1)$ .  
 The standard enthalpy of combustion of 2 mol of benzene is  $-x$  kJ.  
 $x = \underline{\hspace{2cm}}$ .
- (1) standard Enthalpy of formation of 1 mol of  $C_6H_6(1)$ , for the reaction  $6C(\text{graphite}) + 3H_2(g) \rightarrow C_6H_6(1)$  is  $48.5 \text{ kJ mol}^{-1}$ .
- (2) Standard Enthalpy of formation of 1 mol of  $CO_2(g)$ , for the reaction  $C(\text{graphite}) + O_{2(g)} \rightarrow CO_2(g)$  is  $-393.5 \text{ kJ mol}^{-1}$ .
- (3) Standard and Enthalpy of formation of 1 mol of  $H_2O(1)$ , for the reaction  $H_2(g) + \frac{1}{2}O_2(g) \rightarrow H_2O(1)$  is  $-286 \text{ kJ mol}^{-1}$ .

**Ans. (6535)**

**Sol.**  $6C(\text{graphite}) + 3H_2(g) \rightarrow C_6H_6(l); \Delta H = 48.5 \text{ kJ/mol}$

$C(\text{graphite}) + O_2(g) \rightarrow CO_2(g); \Delta H = -393.5 \text{ kJ/mol}$

$H_2^{(g)} + \frac{1}{2}(g) \longrightarrow H_2O(l); \Delta H = -286 \text{ kJ/mol}$

equation  $-(1) \times 1 + (2) \times 6 + (3) \times 3$

$-48.5 - 6 \times 393.5 - 3 \times 286$

$= -3267.5 \text{ kJ for 1 mol}$

$= -6535 \text{ kJ for 2 mol}$

**Ans. 6535 kJ**

82. The fusion of chromite ore with sodium carbonate in the presence of air leads to the formation of products A and B along with the evolution of  $CO_2$ . The sum of spin-only magnetic moment values of A and B is \_\_\_ B.M. (Nearest integer)  
 (Given atomic number : C : 6, Na : 11, O : 8, Fe : 26, Cr : 24]

**Ans. (6)**

**Sol.**  $4FeCr_2O_4 + 8Na_2CO_3 + 7O_2 \rightarrow$   
 $8Na_2CrO_4 + 2Fe_2O_3 + 8CO_2$   
A B

Spin only magnetic moment


For  $Na_2CrO_4$   $\mu_B = 0$

For  $Fe_2O_3$   $\mu_B = 5.9$

sum = 5.9

83. X of ethanamine was subjected to reaction with  $\text{NaNO}_2/\text{HCl}$  followed by hydrolysis to liberate  $\text{N}_2$  and  $\text{HCl}$ . The  $\text{HCl}$  generated was completely neutralised by 0.2 moles of  $\text{NaOH}$ . X is \_\_\_\_ g.

Ans. (9)



84. In an atom, total number of electrons having quantum numbers  $n = 4$ ,  $|m_l| = 1$  and  $m_s = -\frac{1}{2}$  is

Ans. (6)

Sol.  $n = 4$

$\ell$                        $m_\ell$

0                        0

1                        -1, 0, +1

2                        -2, -1, 0, +1, +2, +3

So number of orbital associated with

$n = 4$ ,  $|m_\ell| = 1$  are 6

Now each orbital contain one  $e^-$  with  $m_s = -\frac{1}{2}$

85. Using the given figure, the ratio of  $R_f$  values of sample A and sample C is  $x \times 10^{-2}$ . Value of x is





Fig : Paper chromatography of Samples


Ans. (50)



86. In the Claisen-Schmidt reaction to prepare 351 g of dibenzalacetone using 87 g of acetone, the amount of benzaldehyde required is \_\_\_\_\_ g. (Nearest integer)

Ans. (318)


Sol. Claisen Schmidt reaction



mw of benzaldehyde = 106


$106 \times 3 = 318 \text{ gm}$ . Benzaldehyde is required to give 1.5 mole (or 351 gm) product

87. Consider the following single step reaction in gas phase at constant temperature.



The initial rate of the reaction is recorded as  $r_1$  when the reaction starts with 1.5 atm pressure of A and 0.7 atm pressure of B. After some time, the rate  $r_2$  is recorded when the pressure of C becomes 0.5 atm. The ratio  $r_1 : r_2$  is \_\_\_\_\_  $\times 10^{-1}$ . (Nearest integer)

Ans. (315)



$r_1$     1.5 atm    0.7 atm

$r_2$     0.5 atm    0.2 atm                      0.5 atm

$\therefore r = K [P_A]^2 [P_B]$

$r_1 = K [1.5]^2 [0.7]$


$r_2 = K [0.5]^2 [0.2]$

$\frac{r_1}{r_2} = 9 \times \frac{7}{2} = 31.5 = 315 \times 10^{-1}$




Ans. 315



88. The product © in the following sequence of reactions has \_\_\_\_\_ π bonds.



Ans. (4)



π bonds = 4

89. Considering acetic acid dissociates in water, its dissociation constant is  $6.25 \times 10^{-5}$ . If 5 mL of acetic acid is dissolved in 1 litre water, the solution will freeze at  $-x \times 10^{-2}$  °C, provided pure water freezes at 0 °C.

$x =$  \_\_\_\_\_. (Nearest integer)

Given :  $(K_f)_{\text{water}} = 1.86 \text{ K kg mol}^{-1}$ .


density of acetic acid is  $1.2 \text{ g mol}^{-1}$

molar mass of water =  $18 \text{ g mol}^{-1}$ .

molar mass of acetic acid =  $60 \text{ g mol}^{-1}$ .

density of water =  $1 \text{ g cm}^{-3}$

Acetic acid dissociates as



Ans. (19)


Sol. Mass of  $\text{CH}_3\text{COOH} = V \times d$

$$= 5 \text{ ml} \times 1.2 \text{ g/ml}$$

$$= 6 \text{ gm}$$

$$n_{\text{CH}_3\text{COOH}} = \frac{6}{60} = 0.1 \text{ mol}$$

$$m_{\text{CH}_3\text{COOH}} \approx M_{\text{CH}_3\text{COOH}} = \frac{0.1}{1} = 0.1 \text{ M}$$



C

C - C $\alpha$                       C $\alpha$                       C $\alpha$

$$K_a = \frac{C\alpha^2}{1 - \alpha}$$

$$1 - \alpha \approx 1 \Rightarrow K_a = C\alpha^2$$

$$\alpha = \sqrt{\frac{K_a}{C}} = \sqrt{\frac{6.25 \times 10^{-5}}{0.1}} = 25 \times 10^{-3}$$

$$\text{V.f. (i)} = 1 + \alpha(n - 1) = 1 + \alpha(2 - 1) = 1 + \alpha$$

$$= 1 + 25 \times 10^{-3} = 1.025$$

$$\Delta T_f = iK_f m$$

$$= (1.025)(1.86)(0.1)$$

$$= 0.19$$

$$= 19 \times 10^{-2}$$

90. Number of compounds from the following with zero dipole moment is \_\_\_\_\_.

HF, H<sub>2</sub>, H<sub>2</sub>S, CO<sub>2</sub>, NH<sub>3</sub>, BF<sub>3</sub>, CH<sub>4</sub>, CHCl<sub>3</sub>, SiF<sub>4</sub>, H<sub>2</sub>O, BeF<sub>2</sub>

Ans. (6)

Sol. H<sub>2</sub>, CO<sub>2</sub>, BF<sub>3</sub>, CH<sub>4</sub>, SiF<sub>4</sub>, BeF<sub>2</sub>

are symm. molecule so dipole moment is zero